



Reproductive biology and growth of greater forkbeard *Phycis blennoides* (Brünnich, 1768) in Western Algerian Coasts (Osteichthyes, Gadidae)

Sofiane Med el Amine Benghali^{1&2*}, Salim Mouffok², Ali Kherraz^{1&2}, Zitouni Boutiba²

¹Department of Marine Science and Aquaculture, FSNV-UMAB-Mostaganem, Algeria

²Environmental Surveillance Laboratory, Department of Biology, Faculty of natural sciences and life, University Oran, BP 1524 El Menaouar, Algeria

Article published on June 18, 2014

Key words: gonadosomatic index, hepatosomatic index, first maturity, spawning period.

Abstract

The Great forkbeard, *Phycis blennoides* (Brünnich, 1768), is a species widely distributed throughout the Mediterranean Sea and although it is of great importance for fishing industry, little is known about its reproductive characteristics. This study provides first data on reproductive characteristics of this species in the south western Mediterranean, important for management and stock assessment. For this purpose a total of 461 individuals ranging from 17.5 to 43.5 cm in total length were collected monthly in a period of one year using trammel nets. Sex ratio of males to females was 1:0.91. The estimated length where 50% of analyzed individuals were sexually mature was 24.73 cm for females. The peak value of gonad somatic index was recorded in September and continued throughout the October, indicating the highest spawning activity when the Kn values are low.

*Corresponding Author: Sofiane Med el Amine Benghali ✉ sbenghali@yahoo.com

Introduction

The greater forkbeard, *Phycis blennoides* (Brünnich, 1768) is a common gadoid occurring in the Mediterranean and in the North-eastern Atlantic, from Iceland to Morocco (Tortonese, 1975; Fisher *et al.*, 1987; Davis & Edwards, 1988; Whitehead *et al.*, 1989). It is usually found on muddy or sand bottoms in depths of 100-650 m, but occasionally is captured in shallower depths, close to shore (18 m) (Bello & Rizzi, 1988). The total length at first sexual maturity for males and females are 18-20 cm and 22-23 cm respectively (Gallardo-Cabello & Gual-Frau, 1984). In the Mediterranean, *P. blennoides* spawns from June to August (Gallardo-Cabello & Gual-Frau, 1984).

The estimation of age and growth of fish are essential to the study of biology and population dynamics as well as for the management of fisheries resources, they provide a basis for the knowledge of longevity, age recruitment, age of sexual maturity, spawning periods, migration and mortality. They are also used to study the demographic population structure and dynamics.

Taking into account that length at maturity, fecundity and sex-ratio are some of the most important

parameters in studying reproductive dynamics of gadoids population, this study was carried out by examination of annual changes of the gonado-somatic index (GSI), hepato-somatic index (HSI), condition factor (Kn), asymptotic length L_{∞} , growth coefficient (K) and length-weight relationship in order to evaluate the level of the exploitation in the Algerian coasts.

Therefore, the aim of the present paper was to study the population dynamics, reproduction, condition, age and growth of this species. This is the first paper with a complete analysis of the biology of *P. blennoides* in south-western Mediterranean.

Material and methods

A-Reproduction study

A series of biological samples was conducted on specimens of *P. Blennoides* caught by the commercial trawlers in the sampling area of Oran and Mostaganem (Fig.1). Specimens' total length was measured and both sex and maturity were reported for females which represent the focus of our study. Four different stages were defined (Table 1) according to (ICES, 2007).

Table 1. Different stages of maturity of *P. Blennoides* females

MATURITY STAGES	DESCRIPTION
1. Imature/Resting	Small ovaries, with firm consistence and minimal vascularization, transparent or pink grey, without opaque or hyaline oocytes.
2. Developing/Maturing	Medium or large ovaries, pink or yellow to orange, with vascularization variable, present and obvious. Opaque oocytes present but without hyaline oocytes.
3. Spawning	A-Hydrated-Large ovaries, with firm consistence and vascularization, pink or reddish orange. Opaque and hyaline oocytes present.
4. Post-spawning	Small or medium ovaries, flaccid, dark pink, orange or purple. Opaque and hyaline oocytes absent or residual.

Sex-ratio

After sexing of 461 specimens we found a sampling rate of 51.30% of Males significantly more important than females sex ratio (48.70%) the sex ratio (male: female) is 1:0.91 (Table 2).

Table 2. Percentage of sexes in *P. Blennoides* (*p<0,05).

Sex	Total	Percentage
Females	225	48.70%
Males	236	*51.30%
Total	461	100%

The length abundance curve is shown in Fig. 2. Fig. 3 shows a variation of the percentage of females per month. The female's percentage is dominant during fall and spring season, declining in the winter and summer period.




Fig. 3. Distribution of males and females of *Phycis blennoides* by season. X 2 test results show a prevalence of one sex over the other per sampling month.




Fig. 4. Percentages of different stages of sexual maturity in *Phycis blennoides* females per month.

Results were compared with theoretical ϵ (1.96) at a rate of 95% confidence (Table 3). The calculated value of $\epsilon = 0.87$ is less than the value (1.96) given by the table of the z-score; this finding indicates that males are, on average, significantly larger than females. As regards the sexual maturation of females, different stages of maturation of the gonads during different months of the year are shown in Fig. 4.

Table 3. Different size parameters of males and females of *P. blennoides*

Sex	Males	Females
Total	236	225
X (cm)	23.32	25.13
σ_2 (cm) ²	145.44	781.14
ϵ	0.87	Difference significant




Fig. 5. Monthly trend of GSI and HSI with standard errors in *Phycis blennoides*.

Indices of fish condition

In our study we have used three indexes to determine the spawning period of the species in the study area: the gonado-somatic index (GSI), hepatosomatic index (HSI) and condition index (Kn). These allowed to quantify morphological changes of the specimens and to identify reproduction period by studying the evolution of maturity stages of the ovary.

Hepato-Somatic Index (HSI) and Gonado-Somatic Index (GSI) and condition factor (Kn).

Monthly averages of GSI and HSI calculated from 225 females are plotted in Fig. 5. Only one peak were observed corresponding to the maximum annual spawning period of the population. The highest value

of GSI were found in September and the lowest values in October and November

The highest values of the HSI occurred in July and the lowest fall in November (Fig. 5). Fig. 6 shows the condition factor Kn by seasons in both sexes. The values of Kn resulted overweight, thus revealing breeding events and confirming a rapid maturation occurring from July to September when the values of Kn are very low with irregular variations.




Fig. 6. Condition index (Kn) with standard error according to the season in *Phycis blennoides* females.

Length at maturity

For the statistical method, the L50 point estimated the body size at sexual maturity at 24.73 cm (Fig. 7). All data are combined in Table 4. Our results confirmed values reported for Mediterranean.

Fisheries which differ from those from the North Atlantic where specimens length at maturity is longer than that found in the Mediterranean Sea. Total individuals' length of the monthly samples ranged from a minimum of 17.50 centimeters to a maximum of 43.5 cm. Minimum sizes correspond to females and maximum sizes correspond to males.

B- Growth study

The values of growth parameters were calculated using the software LFDA (subroutine ELEFAN) (Kirkwood *et al.*, 2001). Tables 6 and 7 report values of L∞ (asymptotic length), K (coefficient of growth), to (the theoretical age at which the size is zero), also for Φ (growth index). These values, once estimated for *P. blennoides* specimens, were then replaced in

the equation of Von Bertalanffy. Parameters obtained from the equation of Von Bertalanffy did not differ significantly between the two sexes; but asymptotic length, growth rate and growth index resulted slightly different in males.

Table 4. Summary of first sexual maturity length (L50) of *P. blennoides* females and males from different areas (* only females were studied to determine the size at maturity).

Authors	Area	Males (cm)	Females (cm)
Cohen <i>et al.</i> , 1990	Atlantic	18	33
Rotllant <i>et al.</i> , 2002	Mediterranean	19.32	20
Present work 2014	west Algeria	*	24.73

Table 5. Growth parameters of *Phycis blennoides* females and males.

Sex	Females	Males
Parameters	K L∞ to φ	K L∞ to φ
Result	0.94 41 - 0.70 3.199	0.81 42.14 - 0.71 3.158

Biometric relations observed by analysis of relative growth are shown in Table 6. This relationship indicates an upper bound of allometry (b greater than 3) for females in all months of the year.

We can say that the weight of the species grows faster than the cube of length. Such an allometric relationship was observed also in males. In fact, the upper bound also appeared in the allometry of males. The fitting of a and b ($W_{th} = aL^b$) was employed as input data in stock assessment models.

Table 6. Biometric relation in *Phycis blennoides*.

Sex	$W_{th} = aL^b$	
Females	$W_{th} = 0.004$	$L^{3.159}$
Males	$W_{th} = 0.002$	$L^{3.332}$




Fig.7. Size of first maturity in *Phycis blennoides*.

Discussion

This study presents first data of reproductive characteristics of forkbeard in the south-western Mediterranean Sea and therefore results were compared with other gadiform species common for this area.

Sex-ratio

In the 12 months that lasted our study we observed and analyzed a sample of 461 individuals *Phycis blennoides*. This allows us to see that the males are slightly larger than females. More information was obtained through the study of the sex ratio, according to different parameters and sample size during the year.

We found a change in the rate of femininity with a significant dominance during the fall, which seems to correspond with the period when we recorded the maximum peak of the GSI which could correspond to a strong and early maturation of the ovaries.

The sex ratio showed a predominance of males, the catch rate (48.70%) for females (51.30%) for males, the sex ratio (female: male) is 1:0.91. In fact, this value is not significantly different ($\chi^2 = 0.01$, $df = 11$, $p > 0.05$) of the 1:1 theoretical value. The study of sex ratio depending on the size shows the dominance of male individuals up to size 26 cm, and that beyond this size the proportions of females are dominant, to the size where females reach sexual maturity. These results are in agreement with the different regions of the Mediterranean (Gordon *et al*, 1995. Rotllant *et al*,

2002). This predominance of females in older individuals could be explained by several authors by availability or larger female catchability; is a higher natural mortality in males. These studies showed that the females grow faster than males: in four years (Cohen *et al*, 1990).

Spawning period The GSI is a real coefficient of gonadal maturation. Its increase coincides with gametogenesis while its decrease indicates an active spawning (Lahaye, 1972). Tracking monthly changes GSI allowed us to know the times of sexual activity *P.blennoides* and its breeding season.

We observed the presence of gravid females with a maximum of 0.48 GSI September also, there are resting females in summer and winter; these observations could be explained by the fact that after spawning adult females would regain deep waters. Similarly, males have emissions of their sexual products with a maximum GSI 0.50 in September, this shows that the eggs are laid at the studied species could take place in autumn. Our results are almost similar to those obtained by other authors. They all define clearly the spawning period in autumn; this situation is common to all the Mediterranean coast (Gordon and Duncan, 1985) and in the Ionian Sea (Matarresse *et al.*, 1998). Presence of ripe females indicated that spawning of *Phycis blennoides* occurs during early autumn to early winter. Rotllant *et al.* (2002) investigated population of *Phycis blennoides* in the western Mediterranean Sea. Mature females in their study were found only in autumn.

Parallel to the GSI, we studied the HSI since the energy required for gonad maturity comes from fat reserves stored in the liver. The observation of the temporal evolution of the hepato-somatic report *Phycis blennoides* females showing phases of hepatic synthesis and consumption phases of liver lipids. The largest decline HSI is during the autumn which synchronizes perfectly with the period of mass reproduction in females and coincides with the transfer of liver reserves to the gonad. The results of

the study of the condition index (Kn) in females, set and confirm the spawning period in the range from September to November, as from September, the value of Kn begins to decrease with a minimum in October, which could be explained by a loss of organic matter associated with the laying period.

Length of first sexual maturity

We estimated the size at first maturity (L₅₀) to 24.73 cm. According to Rodriguez - Cabello *et al.* (1998), the size at first maturity of females of the Mediterranean is smaller than that found in the North Atlantic; this difference (also found in males) was explained by suggesting a possible relationship between the maturity of the species and latitude (Lam, 1983). Our findings and conclusions support obtained by various authors mentioned above suggests that the reproductive parameters of *Phycis blennoides* differ from one region to another, probably under the influence of various environmental and geographical parameters, as the passage of Atlantic currents entering the Mediterranean through the Strait Gibraltar rich in organic matter providing an ideal enrichment Algerian deep waters (Cartes *et al.*, 2002), Silva (1986) calculated length at first maturity at 41 cm for females and 36 cm for males in Azorean waters and in this study that length was lower and equal to 24.73 cm for females.

Growth and length-weight relationship

Concerning the study of growth, we used the LFDA software (subroutine ELEFAN), and able to analyze the size frequency of this species which is largely untapped in Algeria, and calculate for the first time the growth parameters in the western basin of Algeria. Model Von Bertalanffy (1938) for linear growth was applied to both sexes. This study revealed that the relative growth rates of each sex decreases with age; they are maximum for small sizes because of the rapid growth of young species compared to older then they fall progressively as the size of the fish approaches the asymptotic size.

Our results are in agreement with many authors who have described the Gadidae in general and in particular *Phycis blennoides* faster growth of females $k = (0.94)$ compared to males (0.81) in the Mediterranean (Cohen *et al.*, 1990 ; Massuti *et al.*, 1996).

The adjustment of growth parameters in Von Bertalanffy model shows that the values differ significantly between the sexes, with higher values of growth coefficient (K) for females. Females grow faster, but reach the maximum size (37 cm) before the males which continue to grow beyond the asymptotic size of females (41cm).

In general, the various aspects discussed on the population dynamics of *Phycis blennoides* (Brünnich 1768) on the west coast of Algeria have shown the same pattern as other areas in the Mediterranean, France (Fancuilli *et al.*, 1979), (Casas and Pineiro, 2000) in the north and north-west of the Iberian Sea. The different authors cited above have also observed this same pattern, but with lower values of growth coefficient (K), especially for females. In any case, the growth parameters estimated (which may overestimate or underestimate) the model parameters used in the study area are consistent with slow growth and high durability (between 10.5-20 years) considered for this Gadidae in the Mediterranean.

The results of the length-weight relationship showed a strong majoring allometry in females and males, that is to say that the weight of the species grows faster than the cube of the length. Adjusted for a and b ($W = alb$) serve as input data in the stock assessment models.

These results were previously observed in other Mediterranean areas (Fancuilli *et al.*, 1979) in the Gulf of Lion in France (Romdhani *et al.*, 2010) in the Tunisian coast.

In the conclusion, this study shows that Great forkbeard has group-synchronous ovarian

development and spawning season from early September to late November. The obtained results from this study are important input data for management and stock assessment of this commercially important fish species.

References

- Alonso-Fernandez A, Dominguez-Petit R, Bao M, Rivas CSaborido-Ray F.** 2008. Spawning pattern and reproductive strategy of female pouting *Trisopterus luscus* (Gadidae) on the Galician shelf of north-western Spain. *Aquatic Living Resources* **21**, 383- 393. doi: 10.1051/alr:2008059
- Beverton RJH, Holt SJ.** 1957. On the dynamics of exploited fish populations. *Fishery investigation, series II.*, **19**, 1-533.
- Bilgin S, Bal H, Tasci B.** 2012. Length based growth estimation and reproduction biology of whiting, *Merlangius merlangus* (Nordman, 1840) in the Southeast Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, **12**, 871-881. doi: 10.4194/1303-2712-v12-4-15
- Cartes JE, Abelló P, Lloris D, Carbonell A, Torres P, Maynou F, Gil De Sola L.** 2002. Feeding guilds of western Mediterranean demersal fish and crustaceans: an analysis based on a spring survey. *Scientia Marina*, **66** (Suppl. 2): 209-220.
- Casas JM, Piñeiro C,** 2000. Growth and age estimation of greater fork-beard (*Phycis blennoides* Brännich, 1768) in the north and northwest of the Iberian Peninsula (ICES Division VIIIc and IXa), *Fisheries Research* **47**(1),19 (2000).
- Cohen DM, Inada T, Iwamoto T, Scialabba N.** 1990 – FAO Species catalogue. Gadiform fishes of the world. *Fisheries Synopsis* **125/10**, 442 pp.
- Edwards, P, Pullin RSV Gartner JA.** 1988. Research and development of integrated crop-livestock-fish farming systems in the tropics. ICLARM Studies and Reviews **16**, 53 pp. International Center for Living Aquatic Resources Management, Manila, Philippines.
- Demestre M, Martin P.** 1993. Optimum exploitation of a demersal resource in the western Mediterranean: the fishery of the deep-water shrimp *Aristeus antennatus* (Risso, 1916). *Scientia Marina* , **57**, 175-182.
- Fanciulli G, Relini Orsi L.** 1979. *Biologia di Phycis blennoides* (Brännich.). *Atti della Società Toscana de Scienze Naturali di Pisa.*, ser B, **86** (suppl.), 383-387.
- Fischer W, Schneider M, Bauchot ML** .1987. Fiches FAO d'identification des espèces pour les besoins de la pêche : Méditerranée et Mer Noire (zone de pêche 37). II Vertébrés (FAO species identification sheets for fishery purposes: Mediterranean and Black Sea (fishing area 37). II Vertebrates). FAO, Rome, pp. 761-1530.
- Gallardo-Cabello M, 1986b.** Analisis de las frecuencias de talla por medio de los métodos de Petersen, Cassie y Bhattacharya, para la determinacion de la edad de la brotola *Phycis blennoides* (Brännich, 1768) en el Mediterráneo Occidental (Pisces: Gadidae). *Instituto de Ciencias del Mar y Limnología Universidad Nacional Autónoma de México . México*, **13**, 187-196.
- Gordon JDM, Duncan JAR.** 1985b. The biology of fish of the Family Moridae in the deep-water of the Rockall Trough. *Journal of the Marine Biological Association of the United Kingdom* **65**, 475-485.
- Gordon JDM, Merrett NR, Haedrich RL.** 1995. Environmental and biological aspects of slope-dwelling fishes of the North Atlantic. pp. 1-26 in Hopper, A.G., editor. *Deep-Water Fisheries of the North Atlantic Oceanic Slope.* Kluwer Academic Publishers, Netherlands

- Gulland JA.** 1983. Fish stock assessment. A manual of basic methods. Ed. John Wiley and Sons, Chichester, UK, FAO/Wilies Series on Food and Agriculture, Vol. 1, 223 pp.
- Gutiérrez-Estrada JC, Pulido-Calvo I, Prenda J.** 2000. Gonadosomatic index estimates of an introduced pumpkinseed (*Lepomis gibbosus*) population in a Mediterranean stream, using computational neural networks. *Aquatic Science* **62**, 350- 363. doi:10.1007/PL00001340.
- Kirkwood GP, Auckland R, Zara JS.** 2001. Length Frequency Distribution Analysis (LFDA), Version 5.0. MRAG Ltd., London, UK.
- Lahaye J.** 1972. Cycles sexuels de quelques poissons plats des côtes bretonnes. *Revue des Travaux de l'Institut des Peches Maritimes* , **36** (2) , 191-207.
- Lam T.** 1983. Environmental influences of gonadal activity in fish. In: Hoar W.S., Randall D.J. & Donalson E.M. (eds.): *Fish Physiology*, 9. Academic Press, New York. London, Part B, pp. 65-116.
- Le Cren ED.** 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, **20**, 201-219.
- Leloup J & Olivereau M.** 1951. Données biométrique comparatives sur la rousette (*Scyllium canicula* L.) de la Manche et de la Méditerranée. *Vie et Milieu*, **2**, 182-209.
- Macer CT.** 1974. The reproductive biology of the horse mackerel *Trachurus trachurus* (L.) in the North Sea and English Channel. *Journal of fish Biology*, **6**, 415-438. doi:10.1111/j.1095-8649.1974.tb04558.x
- Matarrese A, D'Onghia G, Basanisi M, Mastrototaro F.** 1998. Spawning and recruitment of *Phycis blennoides* (Phycidae) from the north-western Ionian Sea (middle-eastern Mediterranean). *Italian Journal of Zoologie*, **65**, 203-209. doi: 10.1080/11250009809386814
- Metin G, Ilkyaz AT, Kinacigil HT.** 2008. Growth, Mortality, and Reproduction of Poor Cood (*Trisopterus minutus* Linn., 1758) in the Central Aegean Sea. *Turkish Journal of Zoology*, **32**, 43-51. doi:10.1017/S0025315410000147
- Murua H, Motos L.** 2006. Reproductive strategy and spawning activity of the European hake *Merluccius merluccius* (L.) in the Bay of Biscay. *Journal of Fish Biology*, **69**, 1288–1303. doi:10.1111/j.1095-8649.2006.01169.x
- Pauly D, Moreau J.** 1997. Méthodes pour l'évaluation des ressources halieutiques. Cépaduès-Éditions, Toulouse, 288 pp.
- Relini G, Orsi-Relini L.** 1987. The decline of red shrimps stocks in the Gulf of Genova. *Investigación Pesquera*, **51**, 245-260.
- Rodríguez-Cabello C, Velasco F, Olasso I.** 1998. Reproductive biology of lesser spotted dogfish *Scylliorhinus canicula* (L., 1758) in the Cantabrian Sea. *Scientia Marina*, **62**, 187-191.
- Romdhani A, Chater I, Ktari MH, Dufour JL, Kélig M.** 2010, Âge et croissance de Phycis blennoides (Brünnich, 1768) des côtes tunisiennes, *Ve Rencontres de l'Ichtyologie en France Paris*, 27-30 mars 2012.
- Rotllant G, Moranta J, Massuti E, Sarda F, Morales-Nin B.** 2002. Reproductive biology of three gadiforms fish species through the Mediterranean deep-sea range (147-1850 m). *Scientia Marina* , **66**, 157-166.
- Šantić M, Pallaoro A, Stagličić N, Mikulandra I, Jardas I.** 2010. Co-variation of gonadosomatic index, condition factor and length-weight relationship

of poor cod, *Trisopterus minutus* (Gadidae), in the Eastern Adriatic Sea. *Cybiurn*, **34**, 279-284.

Silva HM. 1986. Reproduction of the forkbeard *Phycis phycis* (Linnaeus, 1766) in Azorean waters. *ICES CM* 1986/G: 60

Sparre P, Venema SC. 1996. Introduction à l'évaluation des stocks des poissons tropicaux. Première partie: Manuel FAO. Document Technique sur les Pêches, **306** (1), Danida, Rome, 401 pp.

Statsoft INC. 2001. STATISTICA (data analysis software system), version 6, www.statsoft.com.

Sparre P, Venema SC. 1998. Introduction to tropic fish stock assessment. Part 1. Manual. FAO Fisheries Technical Paper. No. 306/1, Rev.2. FAO, Rome, pp. 407.

Tortonese E. 1975. Osteichthyes (Pesci ossei), Parte Seconda. In: Fauna d'Italia XI., Edizioni Calderini, Bologna 636 pp.

Tsikliras AC, Antonopoulou E, Stergiou KI. 2010. Spawning period of Mediterranean marine fishes. Reviews in Fish Biology and Fisheries, **20**, 499-538.

Ursin E. 1967. A mathematical model of some aspects of fish growth, respiration and mortality. *Journal of Fisheries Research Board of Canada*, **24**, 2355-2453.

Von Bertalanffy L. 1938. A quantitative theory of organic growth. *Human Biology*, **10**, 181-213.

Whitehead PJP. 1985. FAO species catalogue. Vol. 7. Clupeoid fishes of the world. Part 1. Chirocentridae, Clupeidae and Pristigasteridae. Rome, Italy. FAO Fish. Synop. 125, Vol. 7, Pt. 1, 303 pp.