



Age and growth of *Mullus surmuletus* (L.) in Mostaganem coast, Northwest Algeria

Ali Kherraz^{1&2}, Amel Kherraz², Sofiane Benghali^{1&2}, Salim Mouffok, Zitouni Boutiba²

¹Department of Marine Science and Aquaculture, Faculty of Natural Sciences and Life, University of Mostaganem, Algeria

²Laboratory Environmental Monitoring Network, Department of Biology, Faculty of Natural Sciences and Life, University Of Oran, Algeria

Article published on August 02, 2014

Key words: *Mullus surmuletus*, Mostaganem, relationship, Von Bertalanffy.

Abstract

The age, growth, of *Mullus surmuletus*, caught in Mostaganem (Northwest Algerian Sea) between January and December 2009, were investigated. Sex ratio female to male was 1:1.60. The total length of females ranged from 12.5 cm to 23 cm and of males from 12.3 cm to 22.5 cm. The length-mass relationship for all individuals can be described by the parameters $a = 0.009$ and $b = 2.934$. Fish aged 0-7 years were present in the samples. The parameters of the Von Bertalanffy growth equation obtained for females were $L_{\infty} = 24.70$ cm, $k = 0.37$ cm / year, and $t_0 = -0.37$ year. For males $L_{\infty} = 25.52$ cm, $k = 0.32$ cm / year, $t_0 = -0.71$ year. Significant differences were found in the growth parameters between males and females.

*Corresponding Author: Ali Kherraz ✉ kherrazali@gmail.com

Introduction

Red mullet *Mullus surmuletus* (Linnaeus 1758) is a demersal marine fish that inhabits sandy and rocky substrata, usually at depths <200 m (Hureau, 1986). *M. surmuletus* is a major target species of Mediterranean demersal fisheries (Reñones *et al.*, 1995; Mehanna, 2009) and encountered in shallower soft bottoms, sea grass beds and rocky bottoms (Lombarte *et al.*, 2000; Bautista-Vega, 2008). This species is benthic carnivores and feed on small invertebrates (Gharbi and Ktari, 1981b; Golani and Galil, 1991; Mehanna, 2009; N'Da, 1992; Labropoulou and Eleftheriou, 1997; Vassilopoulou *et al.*, 2001).

In the Mediterranean, most investigators have studied *Mullus surmuletus* jointly with *M. barbatus*, and reports are available comparing chemical composition (Fernández and Val, 1966), age and growth (Bougis, 1952; Gharbi and Ktari, 1981 a; Andaloro and Giarritta, 1985; Morales-Nin, 1986; 1991; Campillo, 1992; Stergiou *et al.*, 1997; Mehanna, 2009), reproduction (Gharbi and Ktari, 1981b), trophic relationships (Gharbi and Ktari, 1979; Golani and Galil, 1991), and biological and fishing aspects (Sánchez *et al.* 1983; Vassilopoulou and Papaconstantinou 1992; Renõnes *et al.*, 1995; Pajeulo *et al.*, 1997) of both species. In Algeria, only Djabali *et al.*, 1993 and Derbal *et al.*, 2001 have studied some biological aspects of this specie.

The aim of our study is analysing the age and the growth aspects of *M. surmutetus* in Mostaganem (Northwest Algeria). The result of the present work will contribute to the knowledge about the age composition and growth of *M. surmuletus* and also to better understanding of its role in the ecosystem. Consequently, this study will be a step forward to the improvement of the fisheries assessment and management of *M. surmuletus* in this area.

Materials and methods

A total of 838 *M. surmuletus* specimens were collected between January and December 2009 from

monthly samples at depths ranging from 10 to 150 m in Mostaganem coast, Mediterranean Sea, using a commercial bottom trawl net (Fig. 1).

A subsample was taken from monthly samples for the biological examination later in the laboratory, the total lengths (TL) of all fish were measured to the nearest cm and the nearest gram total weight (TW), and the sex of each specimen was determined by examining the gonads macroscopically.

The length-weight relationships were determined according to the allometric equation (Sparre *et al.*, 1989): $W = aL^b$, where W is the total body weight (g), L is the total length (cm), while a and b are constants. Statistical comparison of length-weight relationships between sexes was performed with t-tests (Zar, 1999).

Growth was expressed in terms of the von Bertalanffy equation (Beverton and Holt, 1957): $L_t = L_\infty (1 - e^{-K(t-t_0)})$, where: L_∞ : is the asymptotic total length; L_t : the total length at age t ; K : the growth curvature parameter and t_0 is the theoretical age when fish would have been at zero total length.

Growth parameters were estimated according to the non-linear method by using the software FISAT program package (subroutine ELEFAN). In ELEFAN, data are reconstructed to generate "peaks" and "trough", and the goodness of fit index (R_n) is defined by: $R_n = 10^{ESP/ASP/10}$, where the ASP "Available Sum of Peaks" is computed by adding the "best" value of the available "peaks" and the ESP "Explained Sum of Peaks" is computed by summing all the peaks and troughs "hit" by a growth curve of the form: $L_t = L_\infty (1 - EXP(-K(t - t_0) + St_s + Sto))$ where:

$$St_s = (CK/2\pi) * \sin(2\pi(t-t_s)),$$

$St_o = (CK/2\pi) * \sin(2\pi(t_0-t_s))$, and L_t is the length at time t . (Sparre *et al.*, 1989). Growth parameters obtained for males and females were compared using the multivariate Hotelling's T^2 test (Bernard, 1981). For the sake of comparison, the index of overall

growth performance Φ , proposed by Pauly and Munro, 1984 was used. This test provided an indication of the reliability of age estimates since it had been suggested that phi-prime test values were similar for the same species and genera. The test was based on: $\Phi = \log K + 2 \log L_{\infty}$ (Piñeiro and Sainza, 2003).

Results

The length and weight of the *M. surmuletus* ranged from 12 cm to 23 cm in total length (TL) and from 16 g to 120 g in weight, respectively. Most fish were 14 - 16 cm TL, accounting for 64 % and 42 % of females and males, respectively (Fig. 2).

Of 838 specimens captured, 516 were female and 322 were male. Females measured 12 to 24 cm TL ($\bar{X}=18.21$ cm), males 12.5 to 23.5 cm TL ($\bar{X}=17.69$ cm). The length-weight relationship of *Mullus surmuletus* indicated a positive allometry for female and a negative allometry for male by the following equation: ($W_T = a L_T^b$).

Table 2. Growth parameters for *M.surmuletus* females and males.

Parameters	Females				Males			
	k (cm/yr)	L _∞ (cm)	t ₀ (yr)	Φ	k (cm/yr)	L _∞ (cm)	t ₀ (yr)	Φ
Results	0,37	24,70	- 0,37	2.35	0,32	25,52	- 0,71	2.32

Table 3. Von Bertalanffy Equation.

Von Bertalanffy Equation	
Females	$L_t = 24,70(1-e^{-0,37(t+0,37)})$
Males	$L_t = 25,52(1-e^{-0,32(t+0,71)})$

Discussion

The calculated growth performance index (Φ) in Mostaganem waters was 2.35 for females and 2.32 for males. This finding is in agreement with the considerable similarity between the growth performance indice (Φ) calculated for each sex. In order to compare the growth of the *M. surmuletus* population with others, all available literature data of von Bertalanffy growth parameters and Φ values, including results from the present study are compiled (Table 4).

Females: $W = 0.009 * L_T^{3.02}$, $r^2 = 0.935$, $n = 516$

Males: $W = 0.010 * L_T^{2.98}$, $r^2 = 0.977$, $n = 322$

The analysis by sex showed a significant difference in the *b* coefficient (Table 1).

Table 1. Biometric Relations *M.surmuletus* (Linne, 1758).

W _T = a L _T ^b		
Females	$W_{\infty} = 0,009L_T^{3,02}$	$r^2 = 0.935$
Males	$W_{\infty} = 0,010L_T^{2,98}$	$r^2 = 0.977$

The Von Bertalanffy growth parameters for *M. surmuletus* were estimated as L_∞= 24.70 cm, K = 0.37 cm/year and t₀ = - 0.37 year for females, and L_∞ = 25.52 cm, K = 0.32 cm/year and t₀ = - 0.71 year for males (Table 2). Significant differences were found between the growth of males and females.

The positive allometry of the length weight relationship for the whole population agrees with results of other studies (Table 4 and 5). The differences between females and males, with a small difference in allometric coefficient in females than males, are probably due to the different length distributions of the two sexes. In the size weight relationship, this difference between females and males as a whole means that the portion of the population > 21 cm in length has a very low proportion of males (Fig.2).

Table 4. *Mullus surmuletus* length weight and growth parameters in different areas of Mediterranean Sea (F females; M males; F + M females and males; further abbreviations as in Tables 1 and 2).

	L_{∞} (cm)	k (cm/year)	t_0 (year)	Φ	a	b	Area	Source
F+M	32.52	0.1097	-3.64	2.06	0.0073	3.10	Catalonia	Sánchez <i>et al</i> (1983)
F	29.75	0.49	-0.31				Sicilian Channel	Andaloro and Giarritta (1985)
M	26.25	0.41	-0.23					
F	21.82	0.51	-0.112	2.38	0.1403	3.351	Tunisia	Gharbi and Ktari (1981a)
M	19.87	0.49	-0.025	2.28	0.1443	3.28		
F+M	21.51	0.50	-0.116	2.36				
F	31.90	0.20	-2.60	2.32	0.0095	3.1090	Majorca	Renones (1995)
M	25.54	0.27	-2.45	2.25	0.0104	3.0672		
F+M	31.28	0.21	-2.34	2.31	0.0091	3.1203		
-	31.74	0.47	-0.30	2.67			Egypt	Mehanna (2009)
F	24.70	0.37	-0.37	2.35	0.009	3.02	Algeria	Kherraz <i>et al</i> (2014)
M	25.52	0.32	-0.71	2.32	0.010	2.98		

Table 5. Total length-total weight relationships of *M. surmuletus* reported by various studies.

Author	Region	Sex	Size range (cm)	a	b	effective
Morales-Nin (1991)	Majorca	-	-	0.016	2.91	1092
Campillo (1992)	Gulf of Lion	-	-	0.082	3.00	-
Sanche <i>et al</i> (1995)	Spain	-	-			
Gonçalves <i>et al</i> (1996)	Portugal	-	21.5-38	0.029	3.08	299
Abdallah (2002)	Egypt	-	5.4-20.8	0.011	3.03	122
Valle <i>et al</i> (2003)	West Mediterranean	-	7.7-25.4	0.0097	3.07	146
Mendes <i>et al</i> (2004)	Portugal	-	17-38.2	0.039	3.36	108
Ilhan <i>et al</i> (2009)	Izmir Bay	-	6.6-22.6	0.0083	3.12	192
Mukadder <i>et al</i> (2013)	Saros Bay	♀ ♂	11-26.8 11.8-19.8	0.0075 0.0114	3.16 3.01	184 118
Kherraz <i>et al</i> (2014)	Mostaganem	♀ ♂	12-24 12.5-23.5	0.009 0.010	3.02 2.98	516 322




Fig. 1. Study area (Max Sea-marine software V.10).




Fig. 2. Length-frequency distribution by sex of *M. surmuletus*.

As a whole, growth of *Mullus surmuletus* is fast, with females growing at a slightly slower rate than males. There is no apparent difference in the maximum age between sexes, but females are predominant in Age Classes IV and V. The growth of *Mullus surmuletus* recorded in our study is very similar to that reported for this species in other areas, except in Catalonia where it is lower (Table 4). However, the calculated growth parameters describe the exploited population for the trawl fishery, where the oldest fish were 7 yr of age.

The results of the present work will contribute to the knowledge on age composition, growth of *M. surmuletus* in Mostaganem and also to better understanding of its role in marine ecosystem. This information will help fisheries scientists for future studies on *M. surmuletus* populations and may also help to enforce regulations on commercial fisheries with regard to minimum landing size restrictions for this species.

Acknowledgements

We thank the Algerian Ministry of Higher Education and Scientific Research (MESRS) which funded this experimental study within the framework of CNEPRU project No F01820090008. We are grateful to anonymous reviewers for the constructive criticisms of an earlier draft.

References

- Abdallah M.** 2002. Length-weight relationship of fishes caught by trawl off Alexandria, Egypt. Naga International Center for Living Aquatic Resources Management Quarterly **25 (1)**, 19-20.
- Andaloro F.** 1982. Résumé des paramètres biologiques sur *Mullus surmuletus* de la mer Tyrrhénienne méridionale et de la mer Ionienne septentrionale. FAO Fisheries and Aquaculture Report **266**, 87-88.
- Andaloro, F, Prestipino SG.** 1985. Contribution to the knowledge of the age and growth of striped mullet, *Mullus barbatus* (L., 1758) and red mullet *Mullus surmuletus* (L., 1758) in the Sicilian Channel. FAO Fisheries and Aquaculture Report **336**, 89-92.
- Bautista-Vega A A, Letourneur Y, Harmelin-Vivien M, Salen-Picard C.** 2008. Difference in diet and size-related trophic level in two sympatric fish species, the red mullets *Mullus barbatus* and *Mullus surmuletus*, in the Gulf of Lions (north-west Mediterranean Sea). *Journal of Fish Biology* **73 (10)**, 2402-2420.
- Bernard DR.** 1981. Multivariate analysis as a means of comparing growth in fish. *Canadian Journal of Fisheries and Aquatic Sciences* **38 (2)**, 233-236.
- Beverton R JH, Holt, S J.** 1957. On the Dynamics of Exploited Fish Population. Fish. Invest. Min. Agri. London, **19**, 533 pp.
- Bougis P.** 1952. Recherches biometriques sur les Rougets (*Mullus barbatus* et *Mullus surmuletus* L.1758). archs Zool. Expo.gen. **89(2)**, 57-174.
- Campillo A.** 1992. Les pêcheries françaises de Méditerranée: synthèse des connaissances. Institut Français de Recherche pour l'Exploitation de la Mer, France, 206 pp.

Derbal F, Kara MH. 2001. Inventaire des poissons des côtes de l'est algérien. Rapp. Comm. Int. Mer Médit., **36**, 258.

Djabali F, Brahmi B, Maamass M. 1993. Poissons des côtes algériennes. Pelagos (NS), 1-215.

Fernández R, Val MJ. 1966. Contribucion al estudio biologico- quimico del Salmonete de roca (*Mullus surmuletus* L.) y del de fango (*Mullus barbatus* L.), de M/tlaga. Boln Inst esp Oceanogr **124**:1.

Gharbi H, Ktari MH. 1979. Régime alimentaire des rougets (*Mullus barbatus* Linnaeus, 1758 et *Mullus surmuletus* Linnaeus, 1758) du golfe de Tunis. Bull Inst Océanogr Pêche, Salammbô **6 (1- 4)**, 41-52.

Gharbi H, Ktari H. 1981a. Croissance des rougets en Tunisie. Bulletin National Institute of Oceanography, Peche Salammbô **8**, 5-40.

Gharbi H, Ktari H. 1981b. Biologie de *Mullus barbatus* Linnaeus, 1758 et *Mullus surmuletus* Linnaeus, 1758 (poissons, teleosteens, mullides) des cotes tunisiennes, taille et age de premiere maturite sexuelle, cycle sxuel et coefficient de condition. Bulletin National Institute of Oceanography, Peche. Salammbô **8**, 41-51.

Golani D, Galil B. 1991. Trophic relationship of colonizing and indigenous goatfishes (Mullidae) in the eastern Mediterranean with special emphasis on decapods crustaceans. Hydrobiologia **218**, 27-33.

Hureau JC. 1986. Mullidae. In Fishes of the North-Eastern Atlantic and the Mediterranean. Whitehead, P. J., Bauchot, M. L., Hureau, J. c., Nielsen, J. and E. Tortonese (Eds). Paris; UNESCO, 877-882.

Ilhan DU, Akalin S, Özeydin O, Tosunoglu Z, Gurbet R. 2009. Growth and Reproduction of *Mullus surmuletus* L., 1758 in Aegean Sea. Ege Journal of Fisheries and Aquatic Sciences **26(1)**, 1-5.

Labropoulou M, Eleftheriou A. 1997. The foraging ecology of two pairs of congeneric demersal fish species: importance of morphological characteristics in prey selection. *Journal of Fish Biology* **50(2)**, 324-340.

Linnaeus C.1758. Systema naturae per Regna Tria Nature secundum classes ordinus.

Lombarte A, Recasens L, Gonzáles M, Gil De Sola L. 2000. Spatial segregation of two species of Mullidae (*Mullus surmuletus* and *M. barbatus*) in relation to habitat. Marine Ecology Progress Series **206**, 239-249.

Machias A, Somarakis S, Tsimenides N. 1998. Bathymetric distribution and movements of red mullet *Mullus surmuletus*. *Marine Ecology Progress Series* **166**, 247-257.

Max Sea-marine. 2008. Software V10 3.2.1, copyright informatique et mer 1988, 2008.

Mehanna SF. 2009. Growth, mortality and spawning stock biomass of the striped red mullet *Mullus surmuletus*, in the Egyptian Mediterranean waters. Mediterranean Marine Sciences **10 (2)**, 5-17.

Mendes B, Fonseca P, Campos A. 2004. Weight-length relationships for 46 fish species of the Portuguese west coast. Journal of Applied Ichthyology **20 (5)**, 355-361.

Morales-Nin B. 1986. Age and growth of *Mullus barbatus* and *Mullus surmuletus* from the Catalan Sea. Rapp. Proc. Verb. CIESM **30 (2)**, 232.

Morales-Nin B. 1991. Parametros biologicos del salmonete de roca *Mullus surmuletus* (L. 1758), en Mallorca. Boletin del Instituto Espanol de Oceanografia **7 (2)**, 139-147.

Mukadder A, Ali I. 2013. Age, growth and reproduction of *Mullus surmuletus* (Linnaeus, 1758)

in Saros Bay (Northern Aegean Sea) *J. Black Sea/Mediterranean Environment* **Vol. 19, No. 2**, 217-233 (2013).

N'Da K. 1992. Diet of the red mullet *Mullus surmuletus* (Mullidae) in the northern area of the Bay of Biscay (in French). *Cybius* **16 (2)**, 159-167.

Pajeulo JG, Lorenzo JM, Ramos AG, Mendez-Villamil M. 1997. Biology of red mullet *Mullus surmuletus* (Mullidae) off the Canary Islands, Central-East Atlantic. *South African Journal of Marine Sciences* **18**, 265-272.

Pauly D, Munro JL. 1984. Once more on the comparison of growth in fish and invertebrates. *Fishbyte* **2**, 21-21.

Piñeiro C, Saínza M. 2003. Age estimation, growth and maturity of the European hake, *Merluccius merluccius* (Linnaeus, 1758) from Iberian Atlantic waters. *ICES Journal of Marine Science* **60 (5)**, 1086-1102.

Renones O, Messuti E, Morales-Nin B. 1995. Life history of the red mullet *Mullus surmuletus* from the bottom-trawl fishery off the Island of Majorca (north-west Mediterranean). *Marine Biology* **123 (3)**, 411-419.

Sánchez P, Morales-Nin B, Martín P. 1983. The mullets (*Mullus surmuletus* L. 1758, *Mullus barbatus* L. 1758) of the Catalan coast: biological and fishing aspects (mimeo). International Council of the Exploration of the Sea Comm. Meet (Demersal Fish Comm.) **27**, 1-19.

Sánchez F, Gándara F, Gancedo R. 1995. Atlas de los peces demersales de Galicia y el Cantábrico. Otoño 1991-1993. Publicaciones Especiales, Instituto Español de Oceanografía, Madrid, Spain (20): 100 pp.

Sparre P, Ursin E, Venema SC. 1989. Introduction to tropical fish stock assesment. Part I. Manual FAO Fisheries Technical Paper, 1. Rome FAO, No: **306**, 337 pp.

Valle C, Bayle T J, Ramos AA. 2003. Weight-length relationships for selected fish species of the western Mediterranean Sea. *Journal of Applied Ichthyology* **19 (4)**, 261-262.

Vassilopoulou V, Papaconstantinou C. 1992. Preliminary biological data on the striped red mullet (*Mullus surmeletus*) in the Aegean Sea. *FAO Fisheries and Aquaculture Report* **477**, 85-96.

Zar JH. 1999. *Biostatistical Analysis*, 4th Ed. Prentice Hall, Upper Saddle River, NJ, USA.